

Package ‘double.truncation’

October 13, 2022

Type Package

Title Analysis of Doubly-Truncated Data

Version 1.7

Date 2020-9-8

Author Takeshi Emura, Ya-Hsuan Hu, Chung-Yan Huang

Maintainer Takeshi Emura <takeshiemura@gmail.com>

Description Likelihood-based inference methods with doubly-truncated data are developed under various models.

Nonparametric models are based on Efron and Petrosian (1999) <doi:10.1080/01621459.1999.10474187> and Emura, Konno, and Michimae (2015) <doi:10.1007/s10985-014-9297-5>. Parametric models from the special exponential family (SEF) are based on Hu and Emura (2015) <doi:10.1007/s00180-015-0564-z> and Emura, Hu and Konno (2017) <doi:10.1007/s00362-015-0730-y>.

The parametric location-scale models are based on Dorre et al. (2020) <doi:10.1007/s00180-020-01027-6>.

License GPL-2

Encoding UTF-8

LazyData true

NeedsCompilation no

Repository CRAN

Date/Publication 2020-09-08 08:40:03 UTC

R topics documented:

double.truncation-package	2
NPMLE	3
PMLE.loglogistic	4
PMLE.lognormal	5
PMLE.SEF1.free	6
PMLE.SEF1.negative	8
PMLE.SEF1.positive	9

PMLE.SEF2.negative	11
PMLE.SEF3.free	12
PMLE.SEF3.negative	14
PMLE.SEF3.positive	15
PMLE.Weibull	16
simu.Weibull	18
Index	19

double.truncation-package

Analysis of Doubly-Truncated Data

Description

Likelihood-based inference methods with doubly-truncated data are developed under various models. Nonparametric models are based on Efron and Petrosian (1999) <doi:10.1080/01621459.1999.10474187> and Emura, Konno, and Michimae (2015) <doi:10.1007/s10985-014-9297-5>. Parametric models from the special exponential family (SEF) are based on Hu and Emura (2015) <doi:10.1007/s00180-015-0564-z> and Emura, Hu and Konno (2017) <doi:10.1007/s00362-015-0730-y>. The parametric location-scale models are based on Dorre et al. (2020) <doi:10.1007/s00180-020-01027-6>.

Details

Details are seen from the references.

Author(s)

Takeshi Emura, Ya-Hsuan Hu, Chung-Yan Huang Maintainer: Takeshi Emura <takeshiemura@gmail.com>

References

- Dorre A, Emura T (2019) Analysis of Doubly Truncated Data, An Introduction, JSS Research Series in Statistics, Springer
- Dorre A, Huang CY, Tseng YK, Emura T (2020) Likelihood-based analysis of doubly-truncated data under the location-scale and AFT model, Computation Stat, DOI:10.1007/s00180-020-01027-6
- Efron B, Petrosian V (1999). Nonparametric methods for doubly truncated data. *J Am Stat Assoc* 94: 824-834
- Emura T, Konno Y, Michimae H (2015). Statistical inference based on the nonparametric maximum likelihood estimator under double-truncation. *Lifetime Data Analysis* 21: 397-418
- Emura T, Hu YH, Konno Y (2017) Asymptotic inference for maximum likelihood estimators under the special exponential family with double-truncation, *Stat Pap* 58 (3): 877-909
- Hu YH, Emura T (2015) Maximum likelihood estimation for a special exponential family under random double-truncation, *Computation Stat* 30 (4): 1199-229

NPMLE*Nonparametric inference based on the self-consistency method*

Description

Nonparametric maximum likelihood estimates are computed based on the self-consistency method (Efron and Petrosian 1999). The SE is computed from the asymptotic variance derived in Emura et al. (2015).

Usage

```
NPMLE(u.trunc, y.trunc, v.trunc,epsilon=1e-08)
```

Arguments

u.trunc	lower truncation limit
y.trunc	variable of interest
v.trunc	upper truncation limit
epsilon	error tolerance for the self-consistency algorithm

Details

Details are seen from the references.

Value

f	density
F	cumulative distribution
SE	standard error
convergence	Log-likelihood, and the number of iterations

Author(s)

Takeshi Emura

References

- Efron B, Petrosian V (1999). Nonparametric methods for doubly truncated data. *J Am Stat Assoc* 94: 824-834
- Emura T, Konno Y, Michimae H (2015). Statistical inference based on the nonparametric maximum likelihood estimator under double-truncation. *Lifetime Data Analysis* 21: 397-418
- Dorre A, Emura T (2019) Analysis of Doubly Truncated Data, An Introduction, JSS Research Series in Statistics, Springer

Examples

```
## A data example from Efron and Petrosian (1999) ##
y.trunc=c(0.75, 1.25, 1.50, 1.05, 2.40, 2.50, 2.25)
u.trunc=c(0.4, 0.8, 0.0, 0.3, 1.1, 2.3, 1.3)
v.trunc=c(2.0, 1.8, 2.3, 1.4, 3.0, 3.4, 2.6)
NPMLE(u.trunc,y.trunc,v.trunc)
```

PMLE.loglogistic

Parametric Inference for the log-logistic model

Description

Maximum likelihood estimates and their standard errors (SEs) are computed. Also computed are the likelihood value, AIC, and other quantities.

Usage

```
PMLE.loglogistic(u.trunc, y.trunc, v.trunc, epsilon = 1e-5, D1=2, D2=2, d1=2, d2=2)
```

Arguments

u.trunc	lower truncation limit
y.trunc	variable of interest
v.trunc	upper truncation limit
epsilon	error tolerance for Newton-Raphson
D1	Randomize the intial value if $ mu_h - mu_{h+1} > D1$
D2	Randomize the intial value if $ \sigma_h - \sigma_{h+1} > D2$
d1	$U(-d1, d1)$ is added to the intial value of mu
d2	$U(-d2, d2)$ is added to the intial value of sigma

Details

Details are seen from the references.

Value

eta	estimates
SE	standard errors
convergence	Log-likelihood, degree of freedom, AIC, the number of iterations
Score	score vector at the converged value
Hessian	Hessian matrix at the converged value

Author(s)

Takeshi Emura

References

Dorre A, Huang CY, Tseng YK, Emura T (2020-) Likelihood-based analysis of doubly-truncated data under the location-scale and AFT model, Computation Stat, DOI:10.1007/s00180-020-01027-6

Examples

```
## A data example from Efron and Petrosian (1999) ##
y.trunc=c(0.75, 1.25, 1.50, 1.05, 2.40, 2.50, 2.25)
u.trunc=c(0.4, 0.8, 0.0, 0.3, 1.1, 2.3, 1.3)
v.trunc=c(2.0, 1.8, 2.3, 1.4, 3.0, 3.4, 2.6)
PMLE.loglogistic(u.trunc,y.trunc,v.trunc)
```

PMLE.lognormal

Parametric Inference for the lognormal model

Description

Maximum likelihood estimates and their standard errors (SEs) are computed. Also computed are the likelihood value, AIC, and other quantities.

Usage

```
PMLE.lognormal(u.trunc, y.trunc, v.trunc, epsilon = 1e-5, D1=2, D2=2, d1=2, d2=2)
```

Arguments

u.trunc	lower truncation limit
y.trunc	variable of interest
v.trunc	upper truncation limit
epsilon	error tolerance for Newton-Raphson
D1	Randomize the intial value if mu_h-mu_h+1 >D1
D2	Randomize the intial value if sigma_h-sigma_h+1 >D2
d1	U(-d1,d1) is added to the intial value of mu
d2	U(-d2,d2) is added to the intial value of sigma

Details

Details are seen from the references.

Value

eta	estimates
SE	standard errors
convergence	Log-likelihood, degree of freedom, AIC, the number of iterations
Score	score vector at the converged value
Hessian	Hessian matrix at the converged value

Author(s)

Takeshi Emura

References

Dorre A, Huang CY, Tseng YK, Emura T (2020-) Likelihood-based analysis of doubly-truncated data under the location-scale and AFT model, Computation Stat, DOI:10.1007/s00180-020-01027-6

Examples

```
## A data example from Efron and Petrosian (1999) ##
y.trunc=c(0.75, 1.25, 1.50, 1.05, 2.40, 2.50, 2.25)
u.trunc=c(0.4, 0.8, 0.0, 0.3, 1.1, 2.3, 1.3)
v.trunc=c(2.0, 1.8, 2.3, 1.4, 3.0, 3.4, 2.6)
PMLE.lognormal(u.trunc,y.trunc,v.trunc)
```

PMLE.SEF1.free

Parametric inference for the one-parameter SEF model (free parameter space)

Description

Maximum likelihood estimates and their standard errors (SEs) are computed. Also computed are the likelihood value, AIC, and other quantities.

Usage

```
PMLE.SEF1.free(u.trunc, y.trunc, v.trunc,
tau1 = min(y.trunc), tau2 = max(y.trunc), epsilon = 1e-04)
```

Arguments

u.trunc	lower truncation limit
y.trunc	variable of interest
v.trunc	upper truncation limit
tau1	lower support
tau2	upper support
epsilon	error tolerance for Newton-Raphson

Details

Details are seen from the references.

Value

eta	estimates
SE	standard errors
convergence	Log-likelihood, degree of freedom, AIC, the number of iterations
Score	score at the converged value
Hessian	Hessian at the converged value

Author(s)

Takeshi Emura

References

- Hu YH, Emura T (2015) Maximum likelihood estimation for a special exponential family under random double-truncation, Computation Stat 30 (4): 1199-229
- Emura T, Hu YH, Konno Y (2017) Asymptotic inference for maximum likelihood estimators under the special exponential family with double-truncation, Stat Pap 58 (3): 877-909
- Dorre A, Emura T (2019) Analysis of Doubly Truncated Data, An Introduction, JSS Research Series in Statistics, Springer

Examples

```
### Data generation: see Appendix of Hu and Emura (2015) ####
eta_true=-3
eta_u=-9
eta_v=-1
tau=10
n=300

a=u=v=y=c()

j=1
repeat{
  u1=runif(1,0,1)
  u[j]=tau+(1/eta_u)*log(1-u1)
  u2=runif(1,0,1)
  v[j]=tau+(1/eta_v)*log(1-u2)
  u3=runif(1,0,1)
  y[j]=tau+(1/eta_true)*log(1-u3)
  if(u[j]<=y[j]&&y[j]<=v[j]) a[j]=1 else a[j]=0
  if(sum(a)==n) break
  j=j+1
}
mean(a) ## inclusion probability around 0.5

v.trunc=v[a==1]
u.trunc=u[a==1]
y.trunc=y[a==1]

PMLE.SEF1.free(u.trunc,y.trunc,v.trunc)
```

PMLE.SEF1.negative	<i>Parametric inference for the one-parameter SEF model (negative parameter space)</i>
--------------------	--

Description

Maximum likelihood estimates and their standard errors (SEs) are computed. Also computed are the likelihood value, AIC, and other quantities.

Usage

```
PMLE.SEF1.negative(u.trunc, y.trunc, v.trunc, tau1 = min(y.trunc), epsilon = 1e-04)
```

Arguments

u.trunc	lower truncation limit
y.trunc	variable of interest
v.trunc	upper truncation limit
tau1	lower support
epsilon	error tolerance for Newton-Raphson

Details

Details are seen from the references.

Value

eta	estimates
SE	standard errors
convergence	Log-likelihood, degree of freedom, AIC, the number of iterations
Score	score at the converged value
Hessian	Hessian at the converged value

Author(s)

Takeshi Emura, Ya-Hsuan Hu

References

- Hu YH, Emura T (2015) Maximum likelihood estimation for a special exponential family under random double-truncation, *Computation Stat* 30 (4): 1199-229
- Emura T, Hu YH, Konno Y (2017) Asymptotic inference for maximum likelihood estimators under the special exponential family with double-truncation, *Stat Pap* 58 (3): 877-909
- Dorre A, Emura T (2019) Analysis of Doubly Truncated Data, An Introduction, JSS Research Series in Statistics, Springer

Examples

```
### Data generation: see Appendix of Hu and Emura (2015) ###
eta_true=-3
eta_u=-9
eta_v=-1
tau=10
n=300

a=u=v=y=c()

j=1
repeat{
  u1=runif(1,0,1)
  u[j]=tau+(1/eta_u)*log(1-u1)
  u2=runif(1,0,1)
  v[j]=tau+(1/eta_v)*log(1-u2)
  u3=runif(1,0,1)
  y[j]=tau+(1/eta_true)*log(1-u3)
  if(u[j]<=y[j]&&y[j]<=v[j]) a[j]=1 else a[j]=0
  if(sum(a)==n) break
  j=j+1
}
mean(a) ## inclusion probability around 0.5

v.trunc=v[a==1]
u.trunc=u[a==1]
y.trunc=y[a==1]

PMLE.SEF1.negative(u.trunc,y.trunc,v.trunc)
```

PMLE.SEF1.positive

Parametric Inference for the one-parameter SEF model (positive parameter space)

Description

Maximum likelihood estimates and their standard errors (SEs) are computed. Also computed are the likelihood value, AIC, and other quantities.

Usage

```
PMLE.SEF1.positive(u.trunc, y.trunc, v.trunc, tau2 = max(y.trunc), epsilon = 1e-04)
```

Arguments

u.trunc	lower truncation limit
y.trunc	variable of interest
v.trunc	upper truncation limit
tau2	upper support
epsilon	error tolerance for Newton-Raphson

Details

Details are seen from the references.

Value

eta	estimates
SE	standard errors
convergence	Log-likelihood, degree of freedom, AIC, the number of iterations
Score	score at the converged value
Hessian	Hessian at the converged value

Author(s)

Takeshi Emura, Ya-Hsuan Hu

References

- Hu YH, Emura T (2015) Maximum likelihood estimation for a special exponential family under random double-truncation, Computation Stat 30 (4): 1199-229
- Emura T, Hu YH, Konno Y (2017) Asymptotic inference for maximum likelihood estimators under the special exponential family with double-truncation, Stat Pap 58 (3): 877-909
- Dorre A, Emura T (2019) Analysis of Doubly Truncated Data, An Introduction, JSS Research Series in Statistics, Springer

Examples

```
#### Data generation: Appendix of Hu and Emura (2015)
eta_true=3
eta_u=1
eta_v=9
tau=10
n=300

a=u=v=y=c()

j=1
repeat{
  u1=runif(1,0,1)
  u[j]=tau+(1/eta_u)*log(u1)
  u2=runif(1,0,1)
  v[j]=tau+(1/eta_v)*log(u2)
  u3=runif(1,0,1)
  y[j]=tau+(1/eta_true)*log(u3)
  if(u[j]<=y[j]&&y[j]<=v[j]) a[j]=1 else a[j]=0
  if(sum(a)==n) break
  j=j+1
}
mean(a) ## inclusion probability around 0.5
```

```

v.trunc=v[a==1]
u.trunc=u[a==1]
y.trunc=y[a==1]

PMLE.SEF1.positive(u.trunc,y.trunc,v.trunc)

```

PMLE.SEF2.negative *Parametric Inference for the two-parameter SEF model (negative parameter space for eta_2)*

Description

Maximum likelihood estimates and their standard errors (SEs) are computed. Also computed are the likelihood value, AIC, and other quantities. Since this is the model, estimates for the mean and SD are also computed.

Usage

```
PMLE.SEF2.negative(u.trunc, y.trunc, v.trunc, epsilon = 1e-04)
```

Arguments

u.trunc	lower truncation limit
y.trunc	variable of interest
v.trunc	upper truncation limit
epsilon	error tolerance for Newton-Raphson

Details

Details are seen from the references.

Value

eta	estimates
SE	standard errors
convergence	Log-likelihood, degree of freedom, AIC, the number of iterations
Score	score vector at the converged value
Hessian	Hessian matrix at the converged value

Author(s)

Takeshi Emura, Ya-Hsuan Hu

References

- Hu YH, Emura T (2015) Maximum likelihood estimation for a special exponential family under random double-truncation, *Computation Stat* 30 (4): 1199-229
- Emura T, Hu YH, Konno Y (2017) Asymptotic inference for maximum likelihood estimators under the special exponential family with double-truncation, *Stat Pap* 58 (3): 877-909
- Dorre A, Emura T (2019) Analysis of Doubly Truncated Data, An Introduction, JSS Research Series in Statistics, Springer

Examples

```
### Data generation: see Appendix of Hu and Emura (2015)
n=300
eta1_true=30
eta2_true=-0.5
mu_true=30
mu_u=29.09
mu_v=30.91

a=u=v=y=c()

###generate n samples of (ui,yi,vi) subject to ui<=yi<=vi###
j=1
repeat{
  u[j]=rnorm(1,mu_u,1)
  v[j]=rnorm(1,mu_v,1)
  y[j]=rnorm(1,mu_true,1)
  if(u[j]<=y[j]&&y[j]<=v[j]) a[j]=1 else a[j]=0
  if(sum(a)==n) break ###we need n data set###
  j=j+1
}
mean(a) ### inclusion probability around 0.5 ###

v.trunc=v[a==1]
y.trunc=y[a==1]
u.trunc=u[a==1]

PMLE.SEF2.negative(u.trunc,y.trunc,v.trunc)
```

PMLE.SEF3.free

Parametric Inference for the three-parameter SEF model (free parameter space for eta_3)

Description

Maximum likelihood estimates and their standard errors (SEs) are computed. Also computed are the likelihood value, AIC, and other quantities.

Usage

```
PMLE.SEF3.free(u.trunc, y.trunc, v.trunc,
  tau1 = min(y.trunc), tau2 = max(y.trunc),
  epsilon = 1e-04, D1=20, D2=10, D3=1, d1=6, d2=0.5)
```

Arguments

u.trunc	lower truncation limit
y.trunc	variable of interest
v.trunc	upper truncation limit
tau1	lower support
tau2	upper support
epsilon	error tolerance for Newton-Raphson
D1	Divergence condition for eta_1
D2	Divergence condition of eta_2
D3	Divergence condition of eta_3
d1	Range of randomization for eta_1
d2	Range of randomization for eta_2

Details

Details are seen from the references.

Value

eta	estimates
SE	standard errors
convergence	Log-likelihood, degree of freedom, AIC, the number of iterations
Score	score vector at the converged value
Hessian	Hessian matrix at the converged value

Author(s)

Takeshi Emura, Ya-Hsuan Hu

References

- Hu YH, Emura T (2015) Maximum likelihood estimation for a special exponential family under random double-truncation, *Computation Stat* 30 (4): 1199-229
- Emura T, Hu YH, Konno Y (2017) Asymptotic inference for maximum likelihood estimators under the special exponential family with double-truncation, *Stat Pap* 58 (3): 877-909
- Dorre A, Emura T (2019) Analysis of Doubly Truncated Data, An Introduction, JSS Research Series in Statistics, Springer

Examples

```
## The first 10 samples of the childhood cancer data ##
y.trunc=c(6,7,15,43,85,92,96,104,108,123)
u.trunc=c(-1643,-24,-532,-1508,-691,-1235,-786,-261,-108,-120)
v.trunc=u.trunc+1825
PMLE.SEF3.free(u.trunc,y.trunc,v.trunc)
```

PMLE.SEF3.negative

Parametric Inference for the three-parameter SEF model (negative parameter space for eta_3)

Description

Maximum likelihood estimates and their standard errors (SEs) are computed. Also computed are the likelihood value, AIC, and other quantities.

Usage

```
PMLE.SEF3.negative(u.trunc, y.trunc, v.trunc, tau1 = min(y.trunc),
epsilon = 1e-04, D1=20, D2=10, D3=1, d1=6, d2=0.5)
```

Arguments

u.trunc	lower truncation limit
y.trunc	variable of interest
v.trunc	upper truncation limit
tau1	lower support
epsilon	error tolerance for Newton-Raphson
D1	Divergence condition for eta_1
D2	Divergence condition of eta_2
D3	Divergence condition of eta_3
d1	Range of randomization for eta_1
d2	Range of randomization for eta_2

Details

Details are seen from the references.

Value

eta	estimates
SE	standard errors
convergence	Log-likelihood, degree of freedom, AIC, the number of iterations
Score	score vector at the converged value
Hessian	Hessian matrix at the converged value

Author(s)

Takeshi Emura, Ya-Hsuan Hu

References

- Hu YH, Emura T (2015) Maximum likelihood estimation for a special exponential family under random double-truncation, *Computation Stat* 30 (4): 1199-229
- Emura T, Hu YH, Konno Y (2017) Asymptotic inference for maximum likelihood estimators under the special exponential family with double-truncation, *Stat Pap* 58 (3): 877-909
- Dorre A, Emura T (2019) Analysis of Doubly Truncated Data, An Introduction, JSS Research Series in Statistics, Springer

Examples

```
## The first 10 samples of the childhood cancer data ##
y.trunc=c(6,7,15,43,85,92,96,104,108,123)
u.trunc=c(-1643,-24,-532,-1508,-691,-1235,-786,-261,-108,-120)
v.trunc=u.trunc+1825
PMLE.SEF3.negative(u.trunc,y.trunc,v.trunc)
```

PMLE.SEF3.positive *Parametric Inference for the three-parameter SEF model (positive parameter space for eta_3)*

Description

Maximum likelihood estimates and their standard errors (SEs) are computed. Also computed are the likelihood value, AIC, and other quantities.

Usage

```
PMLE.SEF3.positive(u.trunc, y.trunc, v.trunc, tau2 = max(y.trunc),
epsilon = 1e-04, D1=20, D2=10, D3=1, d1=6, d2=0.5)
```

Arguments

u.trunc	lower truncation limit
y.trunc	variable of interest
v.trunc	upper truncation limit
tau2	upper support
epsilon	error tolerance for Newton-Raphson
D1	Divergence condition for eta_1
D2	Divergence condition of eta_2
D3	Divergence condition of eta_3
d1	Range of randomization for eta_1
d2	Range of randomization for eta_2

Details

Details are seen from the references.

Value

eta	estimates
SE	standard errors
convergence	Log-likelihood, degree of freedom, AIC, the number of iterations
Score	score vector at the converged value
Hessian	Hessian matrix at the converged value

Author(s)

Takeshi Emura, Ya-Hsuan Hu

References

- Hu YH, Emura T (2015) Maximum likelihood estimation for a special exponential family under random double-truncation, *Computation Stat* 30 (4): 1199-229
- Emura T, Hu YH, Konno Y (2017) Asymptotic inference for maximum likelihood estimators under the special exponential family with double-truncation, *Stat Pap* 58 (3): 877-909
- Dorre A, Emura T (2019) Analysis of Doubly Truncated Data, An Introduction, JSS Research Series in Statistics, Springer

Examples

```
## The first 10 samples of the childhood cancer data ##
y.trunc=c(6,7,15,43,85,92,96,104,108,123)
u.trunc=c(-1643,-24,-532,-1508,-691,-1235,-786,-261,-108,-120)
v.trunc=u.trunc+1825
PMLE.SEF3.positive(u.trunc,y.trunc,v.trunc)
```

Description

Maximum likelihood estimates and their standard errors (SEs) are computed. Also computed are the likelihood value, AIC, and other quantities.

Usage

```
PMLE.Weibull(u.trunc, y.trunc, v.trunc, epsilon = 1e-5, D1=2, D2=2, d1=2, d2=2)
```

Arguments

u.trunc	lower truncation limit
y.trunc	variable of interest
v.trunc	upper truncation limit
epsilon	error tolerance for Newton-Raphson
D1	Randomize the intial value if $ \mu_h - \mu_{h+1} > D1$
D2	Randomize the intial value if $ \sigma_h - \sigma_{h+1} > D2$
d1	$U(-d1, d1)$ is added to the intial value of mu
d2	$U(-d2, d2)$ is added to the intial value of sigma

Details

Details are seen from the references.

Value

eta	estimates
SE	standard errors
convergence	Log-likelihood, degree of freedom, AIC, the number of iterations
Score	score vector at the converged value
Hessian	Hessian matrix at the converged value

Author(s)

Takeshi Emura

References

Dorre A, Huang CY, Tseng YK, Emura T (2020) Likelihood-based analysis of doubly-truncated data under the location-scale and AFT model, Computation Stat, DOI:10.1007/s00180-020-01027-6

Examples

```
## A data example from Efron and Petrosian (1999) ##
y.trunc=c(0.75, 1.25, 1.50, 1.05, 2.40, 2.50, 2.25)
u.trunc=c(0.4, 0.8, 0.0, 0.3, 1.1, 2.3, 1.3)
v.trunc=c(2.0, 1.8, 2.3, 1.4, 3.0, 3.4, 2.6)
PMLE.Weibull(u.trunc,y.trunc,v.trunc)
```

simu.Weibull*Simulating doubly-truncated data from the Weibull model***Description**

A data frame is generated by simulated data from the Weibull model.

Usage

```
simu.Weibull(n,mu,sigma,delta)
```

Arguments

n	sample size
mu	location parameter
sigma	scale parameter
delta	a positive parameter controlling the inclusion probability

Details

The data are generated from the random vector (U,Y,V) subject to the inclusion criterion $U \leq Y \leq V$. The random vector are defined as $U = \mu - \delta + \sigma W$, $Y = \mu + \sigma W$, and $V = \mu + \delta + \sigma W$, where $P(W > w) = \exp(-\exp(w))$. See Section 5.1 of Dorre et al. (2020-) for details. The inclusion probability is $P(U \leq Y \leq V)$.

Value

u	lower truncation limits
y	log-transformed lifetimes
v	upper truncation limits

Author(s)

Takeshi Emura

References

Dorre A, Huang CY, Tseng YK, Emura T (2020-) Likelihood-based analysis of doubly-truncated data under the location-scale and AFT model, Computation Stat, DOI:10.1007/s00180-020-01027-6

Examples

```
## A simulation from Dorre et al.(2020) ##
simu.Weibull(n=100,mu=5,sigma=2,delta=2.08)

Dat=simu.Weibull(n=100,mu=5,sigma=2,delta=2.08)
PMLE.Weibull(Dat$u,Dat$y,Dat$v)
```

Index

- * **Exponential distribution**
 - PMLE.SEF1.free, 6
 - PMLE.SEF1.negative, 8
 - PMLE.SEF1.positive, 9
- * **Location-scale family**
 - PMLE.loglogistic, 4
 - PMLE.lognormal, 5
 - PMLE.Weibull, 16
 - simu.Weibull, 18
- * **NPMLE**
 - NPMLE, 3
- * **Normal distribution**
 - PMLE.SEF2.negative, 11
- * **Self-consistency method**
 - NPMLE, 3
- * **Skew normal distribution**
 - PMLE.SEF3.free, 12
 - PMLE.SEF3.negative, 14
 - PMLE.SEF3.positive, 15
- * **Special exponential family**
 - PMLE.SEF1.free, 6
 - PMLE.SEF1.negative, 8
 - PMLE.SEF1.positive, 9
 - PMLE.SEF2.negative, 11
 - PMLE.SEF3.free, 12
 - PMLE.SEF3.negative, 14
 - PMLE.SEF3.positive, 15
- * **Weibull distribution**
 - PMLE.loglogistic, 4
 - PMLE.lognormal, 5
 - PMLE.Weibull, 16
 - simu.Weibull, 18
- * **package**
 - double.truncation-package, 2

double.truncation

- (double.truncation-package), 2

double.truncation-package, 2

NPMLE, 3